

Extended-body approach to the electromagnetic self-force in curved spacetime

Javier Molina Sánchez and Eric Poisson

Department of Physics, University of Guelph, Guelph, Ontario, Canada N1G 2W1

(Dated: May 23, 2006; updated January 5, 2007)

We offer a novel derivation of the electromagnetic self-force acting on a charged particle moving in an arbitrary curved spacetime. Our derivation is based on a generalization from flat spacetime to curved spacetime of the extended-body approach of Ori and Rosenthal. In this approach the charged particle is first modeled as a body of finite extension s , the net force acting on the extended body is computed, and the limit $s \rightarrow 0$ is taken at the end of the calculation. Concretely our extended body is a dumbbell that consists of two point charges that are maintained at a constant spacelike separation s . The net force acting on the dumbbell includes contributions from the mutual forces exerted on each charge by the field created by the other charge, the individual self-forces exerted on each charge by its own field, and the external force which is mostly responsible for the dumbbell's acceleration. These contributions are added up, in a way that respects the curved nature of the spacetime, and all diverging terms in the net force are shown to be removable by mass renormalization. Our end result, in the limit $s \rightarrow 0$, is the standard expression for the electromagnetic self-force in curved spacetime.

Disclaimer

An earlier version of this paper was submitted for publication in *Physical Review D*, and it was severely criticised by a knowledgeable referee. After revising the paper to fully acknowledge the flaws of this work, and failing once more to convince the referee, we decided to withdraw the paper and not to pursue its publication. We tried to do what should not be done: To derive the equations of motion of a charged, extended body without formulating a consistent model for its internal dynamics. This effort was misguided, and the referee was correct to recommend against publication.

Nevertheless, there is much that we like about this paper. First, we believe that Fermi normal coordinates (Sec. II) are a very useful and natural tool for this problem. Second, we find it remarkable that the total force is largely insensitive to the details of the rule (Sec. V) that is used to transport individual forces to a common point. And finally, we believe that our critical review of the literature (Sec. VI) is valuable, and we reiterate our view that “the existing derivations [of the standard expression for the electromagnetic self-force], including the one presented here, possess varying strengths and weaknesses, but that none of them can be considered to be completely satisfactory.” We hope that this contribution will motivate other researchers to seek a derivation that is fully beyond reproach.

I. INTRODUCTION AND OVERVIEW

A particle with an electric charge e , subjected to an external force f_{ext}^μ , moves in a curved spacetime according to the equations of motion

$$ma^\mu = f_{\text{ext}}^\mu + f_{\text{self}}^\mu, \quad (1.1)$$

where m is the particle's mass, $a^\mu = Du^\mu/d\tau$ is the particle's covariant acceleration, the covariant derivative with

respect to proper time τ of the velocity vector u^μ , and f_{self}^μ is the particle's electromagnetic self-force, the force exerted on the particle by its own electromagnetic field. This is given by

$$f_{\text{self}}^\mu = e^2 (\delta^\mu_\nu + u^\mu u_\nu) \left(\frac{2}{3} \frac{Da^\nu}{d\tau} + \frac{1}{3} R^\nu_\lambda u^\lambda \right) + 2e^2 u_\nu \int_{-\infty}^{\tau^-} \nabla^{[\mu} G^{\nu]}_{\lambda'}(z(\tau), z(\tau')) u^{\lambda'} d\tau'. \quad (1.2)$$

The self-force is expressed in terms of the velocity vector u^μ , the rate of change of the acceleration vector $Da^\mu/d\tau$, the Ricci tensor R^μ_ν of the curved spacetime, and in terms of an integral over the particle's past history. The integration involves the retarded Green's function $G^\mu_{\nu'}(x, x')$ associated with the wave equation satisfied by the electromagnetic vector potential (in the Lorenz gauge); the first argument of the Green's function is the point $z(\tau)$, the current position of the particle, and its second argument is a prior position $z(\tau')$. The integration also involves the charge's velocity vector $u^{\mu'}$ at the prior position $z(\tau')$. The integration is cut short at $\tau' = \tau^- \equiv \tau - 0^+$ to avoid the singular behavior of the Green's function when $z(\tau')$ coincides with $z(\tau)$. We shall refer to the right-hand side of Eq. (1.2) as the *standard expression* for the electromagnetic self-force.

The standard expression was first obtained by DeWitt and Brehme [1], who based their derivation on energy-momentum conservation, and who generalized Dirac's pioneering analysis [2] from flat spacetime to curved spacetime; their calculations were later corrected by Hobbs [3]. The standard expression was also derived on the basis of a logically independent axiomatic approach by Quinn and Wald [4]. Another alternative derivation was produced by Detweiler and Whiting [5], on the basis of their unique decomposition of the retarded field into singular and regular fields, and an assertion that the singular field exerts no force on the particle. Finally, the standard expression was derived by Poisson [6] on the basis of an averaging

procedure, wherein the self-force is assumed to be produced by the average of the particle's retarded field over a spherical surface surrounding the charge.

Our purpose in this paper is to propose yet another derivation of the standard expression, and to critically review the derivations listed in the preceding paragraph. Our contention is that the existing derivations, including the one presented here, possess varying strengths and weaknesses, but that none of them can be considered to be completely satisfactory. The derivation presented here is certainly not immune to criticism. We believe, however, that its flaws are sufficiently different from the flaws of the competition that it merits to be developed. Furthermore, we believe that the calculational techniques introduced here will provide a useful basis for a future attempt at providing a fully satisfactory derivation of the standard expression.

We have in mind an approach in which the charged object is fundamentally not a point particle, but a body of finite extension (denoted s) that is held together against electrostatic repulsion by cohesive forces. A physical model for such an object would involve the specification of its internal composition, a description of the internal dynamics (the motion of each body element around the center of mass, in response to the applied cohesive and electromagnetic forces), and a derivation of its external dynamics (the motion of the center of mass in response to the external and self forces). This modeling would have to be done in a fully relativistic setting that takes into account the retardation of all interactions within the body, and the curved nature of the background spacetime. In the limit in which the length and time scales associated with the external electromagnetic field, and those associated with the curved spacetime, are large compared with s , we would expect this detailed calculation to reproduce the standard expression of Eq. (1.2). The reason is that in this limit, the external fields do not probe the internal structure of the charged body, and the equations of motion would have to reflect a near-complete decoupling of the internal and external dynamics; this is achieved by the standard expression.

The program outlined in the previous paragraph is a difficult one to pursue. This approach has been very successful in the context of the self-forced motion of a *massive body* (as opposed to an electrically charged body) in a curved spacetime [6, 7, 8]. Here the body could be modeled as a black hole, an extended object with the simplest structure compatible with the laws of general relativity. The motion of the black hole could be derived by matching the metric of the black hole perturbed by the tidal gravitational field supplied by the external universe to the metric of the external universe perturbed by the moving black hole. Demanding that this metric be a valid solution to the Einstein field equations determines the motion without additional input. The end result is an equation of motion that makes no reference to the internal structure of the extended body. While the gravitational implementation of this program was made

possible by the simplicity afforded by the black hole, an electromagnetic implementation would have to involve a detailed modeling of the body's internal structure. For this reason the electromagnetic case is much more difficult to deal with. To the best of our knowledge, the electromagnetic program has yet to be completed.

In this paper we take some steps toward an eventual completion of this program. We consider a charged body with a very simple internal structure, but we make no attempt to model its internal dynamics. This is hardly satisfactory, and we are indeed a long way from reaching the stated goals, but we nevertheless shall see that some measure of success can be achieved with this minimalistic setting. In particular, we shall see that our approach does permit a derivation of the standard expression of Eq. (1.2). While we concede that our approach is flawed and does not produce a fully satisfactory derivation of Eq. (1.2), we believe that our contribution is still valuable. In particular, as was stated above, we believe that the calculational techniques introduced in this paper will be useful in an eventual completion of the program.

Concretely our derivation is based on the Ori-Rosenthal extended-body approach to the electromagnetic self-force [9, 10], which we generalize from flat spacetime to curved spacetime. Our extended body is a dumbbell that consists of two point charges that are maintained at a constant spacelike separation s ; the orientation of the dumbbell, relative to the direction of the motion, is arbitrary. The dumbbell has no internal dynamics: the separation between the two charges is kept constant by a method that will be described in Sec. IV. Because our extended body does not have a continuous distribution of charge, our calculation of the net force acting on it cannot proceed without an input axiom; this must provide us with some information on the individual self-forces that act on the constituting charges. We want, of course, this axiom to be as mild as possible, and we want the calculation to be capable of producing, in the limit $s \rightarrow 0$, a precise and well-defined expression for the electromagnetic self-force.

Our input axiom is mild. It states:

The self-force acting on a point charge is a well-defined vector field on the particle's world line, and it is proportional to the square of the particle's electric charge.

This axiom is, first of all, an assumption that there exists a self-force acting on each constituting point charge in the dumbbell. This assumption is required by physical consistency: It would be inconsistent to assume that the individual self-forces do not exist, and to find in the end that a nonzero self-force emerges from the extended body in the limit $s \rightarrow 0$. The axiom is also an assumption that the self-force is proportional to the square of the particle's charge. This property also follows from a requirement of physical consistency: In a spacetime with a timelike Killing vector, the total work done by the self-force must equal the total energy radiated by the charge

[11]; because the radiated energy is proportional to the square of the charge, we require the same of the self-force.

The formulation of the axiom constitutes the first flaw of our minimalistic approach: such an axiom would not be required in an extended-body calculation that would deal honestly with the body's internal dynamics. Moreover, the assumed equality between radiated energy and work done by the self-force must be only approximately valid, because heat dissipation within an extended body would also contribute to the energy balance. This effect, however, should become unimportant in the limit $s \rightarrow 0$, that is, in the limit in which the internal dynamics decouple from the external dynamics.

Our derivation of the standard expression rests on the axiom, on additional contrivances that will be introduced below, and on various computational devices that can be imported from the literature. The geometry of our dumbbell is well adapted to a description in terms of Fermi normal coordinates, and we make heavy use of this coordinate system throughout the paper. The coordinates are introduced in most textbooks on general relativity (see, for example, Refs. [12] or [13]), but here we rely on the presentation given in Poisson's review article, published online in *Living Reviews in Relativity* [6]. We will make heavy use of this resource in this paper; below it will be repeatedly referred to as LRR. The properties of the Fermi normal coordinates that are needed in our derivation are summarized in Sec. II.

Another set of results that can be imported from the literature is the expression for the retarded electromagnetic field created by a point charge moving on an accelerated world line in a curved spacetime. This field produces a force on a neighboring charge, and this force contributes to the net force acting on the dumbbell. Expressions for the field, given as expansions in powers of s in Fermi normal coordinates, are derived in LRR [6] and used here; these are presented in Sec. III.

A useful computational device for our derivation is the Detweiler-Whiting decomposition [5] of the retarded electromagnetic field $F_{\alpha\beta}$ into singular "S" and regular "R" pieces,

$$F_{\alpha\beta} = F_{\alpha\beta}^S + F_{\alpha\beta}^R. \quad (1.3)$$

As was shown by Detweiler and Whiting, this decomposition is unambiguous, the retarded and singular fields share the same singularity structure near the world line, and the regular field $F_{\alpha\beta}^R$ is smooth on the world line. They showed also that the retarded and singular fields satisfy the same field equations (with a distributional current density on the right-hand side), but that the regular field is sourcefree. The Detweiler-Whiting decomposition is introduced in Sec. III, and it is used here without making assumptions about the physical role of the singular field. We involve it mostly as a matter of convenience, but also to emphasize the point that the standard expression for the self-force describes an interaction between the particle and the regular field.

The dumbbell is formally introduced in Sec. IV, and

in Sec. V the net force acting on it is computed from the individual contributions. These include the mutual forces exerted on each charge by the field created by the other charge, the individual self-forces exerted on each charge by its own field, and the external force which is mostly responsible for the dumbbell's acceleration. This step requires another computational device, a transport rule that allows us to bring the different forces to a common point before adding them up. Our transport rule is introduced in Sec. V, and it is a straightforward generalization, from flat spacetime to curved spacetime, of the Ori-Rosenthal rule [9, 10].

The final step of the calculation is to renormalize the mass and then take the limit $s \rightarrow 0$. These calculations also are presented in Sec. V, and the final outcome is Eq. (1.2), the standard expression for the electromagnetic self-force. We conclude the paper in Sec. VI with a critical review of the existing derivations of the self-force, in which we comment on their strengths and weaknesses. We also offer a critical summary of our own approach.

Throughout the paper we use the notations and conventions of Misner, Thorne, and Wheeler [12]. In particular, we use geometrized units in which $G = c = 1$.

II. FERMI COORDINATES AND COMOVING WORLD LINES

A. Fermi normal coordinates

We consider a neighborhood $\mathcal{N}$ around a reference world line $\bar{\gamma}$ in a curved spacetime with metric $g_{\alpha\beta}$; the extension of $\mathcal{N}$ is small compared with the radius of curvature of the spacetime. The world line is arbitrary, and it possesses an acceleration. The neighborhood $\mathcal{N}$ is charted with Fermi normal coordinates (t, x^a) , in which t is proper time on the world line, and x^a ($a = 1, 2, 3$) are spatial coordinates that vanish on the world line [LRR Sec. 3.2]. The coordinates have the property that a space-like geodesic orthogonal to $\bar{\gamma}$ is described by the equations $t = \text{constant}$ and $x^a = s\omega^a$, where s is proper distance away from the world line (measured along the geodesic), and the coefficients ω^a are constant. They also have the property that $s^2 = \delta_{ab}x^ax^b$, where δ_{ab} is the Kronecker delta, so that the ω^a 's form the components of a unit vector, $\delta_{ab}\omega^a\omega^b = 1$.

The metric of the curved spacetime in $\mathcal{N}$ is expressed in Fermi normal coordinates (FNC) as an expansion in powers of the spatial coordinates x^a . It takes the form [LRR Eq. (125)–(127)]

$$g_{tt} = -1 - 2\bar{a}_a x^a - (\bar{a}_a x^a)^2 - \bar{R}_{ctdt} x^c x^d + O(s^3), \quad (2.1)$$

$$g_{ta} = -\frac{2}{3}\bar{R}_{actd} x^c x^d + O(s^3), \quad (2.2)$$

$$g_{ab} = \delta_{ab} - \frac{1}{3}\bar{R}_{acbd} x^c x^d + O(s^3), \quad (2.3)$$

where $\bar{a}_a(t) \equiv \delta_{ab}\bar{a}^b(t)$ are the spatial components of the vector $a_{\bar{a}}$, $\bar{\gamma}$'s acceleration vector, in FNC (the time

component vanishes), and where $\bar{R}_{ctdt}(t)$, $\bar{R}_{actd}(t)$, and $\bar{R}_{acbd}(t)$ are the components of the Riemann tensor evaluated on $\bar{\gamma}$; these depend on t only. Here and below we use an overbar to indicate quantities that are evaluated on $\bar{\gamma}$. When the quantity is the component of a tensor in FNC, we place the overbar on the tensorial symbol, as in $\bar{a}_a$; when the quantity is the tensor itself, we place the overbar on the tensorial index, as in $a_{\bar{a}}$.

The FNC have the property that the metric reduces to Minkowski values on the reference world line. The inverse metric is

$$g^{tt} = -1 + 2\bar{a}_a x^a - 3(\bar{a}_a x^a)^2 + \bar{R}_{ctdt} x^c x^d + O(s^3), \quad (2.4)$$

$$g^{ta} = -\frac{2}{3}\bar{R}_{ctd}^a x^c x^d + O(s^3), \quad (2.5)$$

$$g^{ab} = \delta^{ab} + \frac{1}{3}\bar{R}_{cd}^a x^c x^d + O(s^3). \quad (2.6)$$

Here the indices on the Riemann tensor are raised freely with δ^{ab} ; this is appropriate because the Riemann tensor is evaluated on the reference world line.

It is useful to decompose the metric in terms of a tetrad of dual vectors $e_{\alpha}^{(0)}$, $e_{\alpha}^{(a)}$ that are orthonormal everywhere in $\mathcal{N}$. Here the superscript within brackets serves to label each one of the four tetrad members, and the Greek subscript is the usual vectorial index.¹ The decomposition is given by

$$g_{\alpha\beta} = -e_{\alpha}^{(0)} e_{\beta}^{(0)} + \delta_{ab} e_{\alpha}^{(a)} e_{\beta}^{(b)} \quad (2.7)$$

and the members of the tetrad are [LRR Eqs. (123) and (124)]

$$e_t^{(0)} = 1 + \bar{a}_a x^a + \frac{1}{2}\bar{R}_{ctdt} x^c x^d + O(s^3), \quad (2.8)$$

$$e_a^{(0)} = \frac{1}{6}\bar{R}_{actd} x^c x^d + O(s^3), \quad (2.9)$$

$$e_t^{(a)} = -\frac{1}{2}\bar{R}_{ctd}^a x^c x^d + O(s^3), \quad (2.10)$$

$$e_b^{(a)} = \delta_b^a - \frac{1}{6}\bar{R}_{cbd}^a x^c x^d + O(s^3). \quad (2.11)$$

The tetrad vectors possess the important property that they are parallel transported along the spacelike geodesic that leaves $\bar{\gamma}$ orthogonally to arrive at the point $x = (t, x^a)$. As was mentioned previously, this spacelike geodesic is described by the equations $t = \text{constant}$ and $x^a = s\omega^a$, with constant coefficients ω^a ; its departure point on $\bar{\gamma}$ is therefore $\bar{x} = (t, 0)$.

We introduce also a tetrad of vectors $e_{(0)}^{\alpha}$, $e_{(a)}^{\alpha}$ which is related to the dual tetrad by

$$e_{(0)}^{\alpha} = -g^{\alpha\beta} e_{\beta}^{(0)}, \quad e_{(a)}^{\alpha} = \delta_{ab} g^{\alpha\beta} e_{\beta}^{(b)}. \quad (2.12)$$

With Eqs. (2.4)–(2.11) we obtain the explicit forms

$$e_{(0)}^t = 1 - \bar{a}_a x^a + (\bar{a}_a x^a)^2 - \frac{1}{2}\bar{R}_{ctdt} x^c x^d + O(s^3), \quad (2.13)$$

$$e_{(0)}^a = \frac{1}{2}\bar{R}_{ctd}^a x^c x^d + O(s^3), \quad (2.14)$$

$$e_{(a)}^t = -\frac{1}{6}\bar{R}_{actd} x^c x^d + O(s^3), \quad (2.15)$$

$$e_{(a)}^b = \delta_b^a + \frac{1}{6}\bar{R}_{cad}^b x^c x^d + O(s^3). \quad (2.16)$$

Any vector A^{α} at a point $x = (t, x^a)$ in $\mathcal{N}$ can be decomposed in the tetrad $e_{(0)}^{\alpha}$, $e_{(a)}^{\alpha}$; the decomposition takes the form of

$$A^{\alpha} = A^{(0)} e_{(0)}^{\alpha} + A^{(a)} e_{(a)}^{\alpha}, \quad (2.17)$$

and the coefficients $A^{(0)} = A^{\alpha} e_{\alpha}^{(0)}$, $A^{(a)} = A^{\alpha} e_{\alpha}^{(a)}$ are the frame components of the vector at the point x .

Any vector A^{α} at the point $x = (t, x^a)$ can be parallel transported to the simultaneous point $\bar{x} = (t, 0)$ on $\bar{\gamma}$ by following the spacelike geodesic $t = \text{constant}$ that links these points. Because the tetrad vectors are themselves parallel transported along this geodesic, the frame components of the vector stay constant during the transport. The vector at $\bar{x}$ is therefore $A^{\bar{\alpha}} = A^{(0)} e_{(0)}^{\bar{\alpha}} + A^{(a)} e_{(a)}^{\bar{\alpha}}$, where $e_{(0)}^{\bar{\alpha}}$ and $e_{(a)}^{\bar{\alpha}}$ are the tetrad vectors at $\bar{x}$. This can be expressed as

$$A^{\bar{\alpha}}(\bar{x}) = g^{\bar{\alpha}}_{\beta}(\bar{x}, x) A^{\beta}(x), \quad (2.18)$$

where $g^{\bar{\alpha}}_{\beta} = e_{(0)}^{\bar{\alpha}} e_{\beta}^{(0)} + e_{(a)}^{\bar{\alpha}} e_{\beta}^{(a)}$ is the parallel propagator. Its components in FNC are

$$g^{\bar{t}}_t = 1 + \bar{a}_a x^a + \frac{1}{2}\bar{R}_{ctdt} x^c x^d + O(s^3), \quad (2.19)$$

$$g^{\bar{t}}_a = \frac{1}{6}\bar{R}_{actd} x^c x^d + O(s^3), \quad (2.20)$$

$$g^{\bar{a}}_t = -\frac{1}{2}\bar{R}_{ctd}^a x^c x^d + O(s^3), \quad (2.21)$$

$$g^{\bar{a}}_b = \delta_b^a - \frac{1}{6}\bar{R}_{cbd}^a x^c x^d + O(s^3). \quad (2.22)$$

B. Congruence of comoving world lines

We introduce a congruence of comoving world lines $\gamma(x^a)$ in $\mathcal{N}$. These timelike curves are all described by $x^a = \text{constant}$, and they are comoving with $\bar{\gamma} \equiv \gamma(0)$ in the sense that they preserve their spatial coordinates, and therefore their spatial separation s relative to $\bar{\gamma}$, during their motion. The comoving world lines are labeled by x^a and are parameterized by their proper time $\tau(x^a)$. This is related to $t \equiv \bar{\tau} \equiv \tau(0)$ by $d\tau^2 = -g_{tt} dt^2$, so that

$$\frac{d\tau}{dt} = 1 + \bar{a}_a x^a + \frac{1}{2}\bar{R}_{ctdt} x^c x^d + O(s^3) \quad (2.23)$$

¹ The dual tetrad is denoted $\bar{e}_{\alpha}^0$, $\bar{e}_{\alpha}^a$ in LRR.

according to Eq. (2.1). The inverse relation is

$$\frac{dt}{d\tau} = 1 - \bar{a}_a x^a + (\bar{a}_a x^a)^2 - \frac{1}{2} \bar{R}_{ctdt} x^c x^d + O(s^3). \quad (2.24)$$

The congruence's velocity field is denoted u^α , and its components in FNC are

$$u^t = \frac{dt}{d\tau}, \quad u^a = 0. \quad (2.25)$$

This vector field is everywhere tangent to the members of the congruence. It is decomposed in the tetrad of Eqs. (2.13)–(2.16) as

$$u^\alpha = u^{(0)} e_{(0)}^\alpha + u^{(a)} e_{(a)}^\alpha, \quad (2.26)$$

and the frame components are calculated to be

$$u^{(0)} = 1 + O(s^3), \quad u^{(a)} = -\frac{1}{2} \bar{R}_{ctd}^a x^c x^d + O(s^3). \quad (2.27)$$

The acceleration field of the congruence is defined as $a^\alpha = u^\beta \nabla_\beta u^\alpha$; this is the acceleration vector of each comoving world line. A straightforward calculation, based on the Christoffel symbols obtained from Eqs. (2.1)–(2.3), produces its components in FNC. We obtain

$$a^t = O(s^2), \quad a^a = (1 - \bar{a}_c x^c) \bar{a}^a + \bar{R}_{tct}^a x^c + O(s^2); \quad (2.28)$$

as expected, $a^t = 0$ and $a^a = \bar{a}^a$ when $x^c = 0$. The acceleration field also can be decomposed as

$$a^\alpha = a^{(0)} e_{(0)}^\alpha + a^{(a)} e_{(a)}^\alpha. \quad (2.29)$$

Its frame components are

$$a^{(0)} = O(s^2), \quad a^{(a)} = (1 - \bar{a}_c x^c) \bar{a}^a + \bar{R}_{tct}^a x^c + O(s^2). \quad (2.30)$$

We conclude this section by introducing one additional vector field in $\mathcal{N}$. Take the spatial coordinates x^a of the point x and divide by the proper distance $s = (\delta_{ab} x^a x^b)^{1/2}$ between x and $\bar{x}$; this gives $\omega^a = x^a/s$, and these form the components of a unit vector. Then define the vector field

$$\omega^\alpha = \omega^a e_{(a)}^\alpha. \quad (2.31)$$

This is well-defined provided that $x^a \neq 0$; the vector field is singular on $\bar{\gamma}$ by virtue of its directional ambiguity. It is easy to see that this vector is normalized, in the sense that $g_{\alpha\beta} \omega^\alpha \omega^\beta = 1$. Its most important property is that it is tangent to the spacelike geodesic that links the points x and $\bar{x}$. This can be established from the definition of the FNC in terms of Synge's world function $\sigma(x, \bar{x})$ [LRR Secs. 2.1 and 3.2, especially Eq. (116)]: We have $x^a = -e_{\bar{\alpha}}^{(a)} \sigma^{\bar{\alpha}}$, where $e_{\bar{\alpha}}^{(a)}$ is the dual tetrad at $\bar{x}$ and $\sigma^{\bar{\alpha}} = \nabla^{\bar{\alpha}} \sigma(x, \bar{x})$, and it follows that

$$\omega^\alpha = -\frac{1}{s} e_{(a)}^\alpha e_{\bar{\alpha}}^{(a)} \sigma^{\bar{\alpha}} = -\frac{1}{s} g_{\bar{\alpha}}^\alpha(x, \bar{x}) \sigma^{\bar{\alpha}},$$

where we have involved the definition of the parallel propagator that follows Eq. (2.18) and used the fact that $e_{\bar{\alpha}}^{(0)} \sigma^{\bar{\alpha}} = -u_{\bar{\alpha}} \sigma^{\bar{\alpha}} = 0$. Using now the identity $-g_{\bar{\alpha}}^\alpha \sigma^{\bar{\alpha}} = \sigma^\alpha$, where $\sigma^\alpha = \nabla^\alpha \sigma(x, \bar{x})$, we arrive at

$$\omega^\alpha = \frac{1}{s} \sigma^\alpha. \quad (2.32)$$

The vector σ^α/s is normalized and it does have the property of being tangent to the geodesic linking x to $\bar{x}$ [LRR Sec. 2.1]. We note that the vector of Eq. (2.32) is defined in a covariant way; the spatial distance s is an invariant related to Synge's world function by $s = \sqrt{2\sigma(x, \bar{x})}$.

III. FORCE ON A COMOVING CHARGE EXERTED BY ANOTHER COMOVING CHARGE

We place an electric charge $\bar{e}$ on the reference world line $\bar{\gamma}$, and we let it create a retarded electromagnetic field $F_{\alpha\beta}$. We next place an electric charge e on the world line $\gamma \equiv \gamma(s\omega^a)$, a specific member of the congruence introduced in Sec. II B. Our specific world line is labeled by its Fermi normal coordinates (FNC) $x^a = s\omega^a$, where s is γ 's constant spatial separation relative to $\bar{\gamma}$, and ω^a are constant directional coefficients. The electromagnetic field created by $\bar{e}$ exerts a force on e , and this is given by

$$f^\alpha = e F_{\alpha\beta} u^\beta = f^{(0)} e_{(0)}^\alpha + f^{(a)} e_{(a)}^\alpha. \quad (3.1)$$

Here u^α is the velocity vector of the charge e moving on the world line γ , whose components are listed in Eqs. (2.24) and (2.25). In the second form the force is decomposed in the tetrad of Eqs. (2.13)–(2.16). In these expressions we must substitute $x^a = s\omega^a$ to select our particular member γ from the congruence $\gamma(x^a)$.

If we further decompose $F_{\alpha\beta}$ and u^α in the tetrad $e_{(0)}^\alpha$, $e_{(a)}^\alpha$ we find that the frame components of the force are given by

$$f^{(0)} = e F_{(a)(0)} u^{(a)}, \quad f^{(a)} = e (F_{(0)}^{(a)} u^{(0)} + F_{(b)}^{(a)} u^{(b)}). \quad (3.2)$$

The frame components of the electromagnetic field were computed, and expressed as expansions in powers of s in FNC, in [LRR Eqs. (455) and (456)]. They are given by²

$$\begin{aligned} F_{(a)(0)} = & \bar{e} \left[\frac{1}{s^2} \omega_a - \frac{1}{2s} (\bar{a}_a + \bar{a}_b \omega^b \omega_a) + \frac{3}{4} \bar{a}_b \omega^b \bar{a}_a \right. \\ & + \frac{3}{8} (\bar{a}_b \omega^b)^2 \omega_a + \frac{3}{8} \bar{a}_t \omega_a - \frac{2}{3} \bar{R}_{atbt} \omega^b \\ & \left. - \frac{1}{6} \bar{R}_{btct} \omega^b \omega^c \omega_a \right] \end{aligned}$$

² The frame components are denoted $\bar{F}_{a0}$ and $\bar{F}_{ab}$ in LRR, and contrary to our usage here, quantities that refer to the reference world line $\bar{\gamma}$ are presented in LRR without an overbar.

$$+ \frac{1}{12}(5\bar{R}_{tt} + \bar{R}_{bc}\omega^b\omega^c + \bar{R})\omega_a - \frac{1}{6}\bar{R}_{ab}\omega^b \Big] + F_{(a)(0)}^R + O(s) \quad (3.3)$$

and

$$F_{(a)(b)} = \bar{e} \left[\frac{1}{2}(\omega_a \dot{a}_b - \dot{a}_a \omega_b) + \frac{1}{2}(\bar{R}_{atbc} - \bar{R}_{btac})\omega^c - \frac{1}{2}(\bar{R}_{at}\omega_b - \omega_a \bar{R}_{bt}) \right] + F_{(a)(b)}^R + O(s). \quad (3.4)$$

The quantities that appear in Eqs. (3.3) and (3.4) were, for the most part, introduced in Sec. II. Exceptions are $\dot{a}_t$ and $\dot{a}_a$, which are the components in FNC of the vector $\dot{a}_{\bar{\alpha}} = Da_{\bar{\alpha}}/d\bar{\tau}$, the covariant derivative with respect to proper time $\bar{\tau} \equiv \tau(0) \equiv t$ of $\bar{\gamma}$'s acceleration vector. Additional exceptions are $\bar{R}_{tt}$, $\bar{R}_{ta}$, $\bar{R}_{ab}$, and $\bar{R}$, which are respectively the components in FNC of the Ricci tensor and the Ricci scalar, all evaluated on the reference world line $\bar{\gamma}$. It should be noted that in Eqs. (3.3) and (3.4), $\omega_a \equiv \delta_{ab}\omega^b$.

We have also introduced the frame components of the Detweiler-Whiting regular field [Eq. (1.3); LRR Eqs. (471) and (472)],

$$F_{(a)(0)}^R = \bar{e} \left[\frac{2}{3}\dot{a}_a + \frac{1}{3}\bar{R}_{at} \right] + \bar{F}_{(a)(0)}^{\text{tail}} + O(s) \quad (3.5)$$

and

$$F_{(a)(b)}^R = \bar{F}_{(a)(b)}^{\text{tail}} + O(s). \quad (3.6)$$

These are related to the frame components of the “tail part” of the field [LRR Eqs. (453) and (454)],

$$\bar{F}_{(a)(0)}^{\text{tail}} = F_{\bar{\alpha}\bar{\beta}}^{\text{tail}} e_{(a)}^{\bar{\alpha}} e_{(0)}^{\bar{\beta}}, \quad \bar{F}_{(a)(b)}^{\text{tail}} = F_{\bar{\alpha}\bar{\beta}}^{\text{tail}} e_{(a)}^{\bar{\alpha}} e_{(b)}^{\bar{\beta}}, \quad (3.7)$$

where

$$F_{\bar{\alpha}\bar{\beta}}^{\text{tail}}(\bar{x}) = 2e \int_{-\infty}^{\bar{\tau}^-} \nabla_{[\bar{\alpha}} G_{\bar{\beta}]\mu'}(\bar{x}, z') u^{\mu'} d\bar{\tau}'. \quad (3.8)$$

Notice that the tail field is evaluated at the point $\bar{x} = (t, 0)$ on the reference world line. It is expressed as an integral over $\bar{e}$'s past history, for times ranging from $t' = -\infty$ to $t' = t^- \equiv t - 0^+$. The first argument of the Green's function is the point $\bar{x}$, the current position of the charge $\bar{e}$, and its second argument is a prior position $z' = (t', 0)$. The tail field is presented in a covariant form by identifying $\bar{\tau}' \equiv \tau'(0) \equiv t'$ and $\bar{\tau} \equiv t$.

If we substitute Eqs. (2.27), (3.3), and (3.4) into Eq. (3.2) we obtain the frame components of the force exerted on the charge e . We find

$$f^{(0)} = O(s), \quad f^{(a)} = eF_{(0)}^{(a)} + O(s). \quad (3.9)$$

We may now substitute these into Eq. (3.1) and reconstruct a vectorial expression for the force. The vector f^α

is at first expressed in terms of the vectors $\omega^a e_{(a)}^\alpha$, $\bar{a}^a e_{(a)}^\alpha$, and quantities of the form $e_{(a)}^\alpha \bar{R}_{tct}^a \omega^c$ and $e_{(a)}^\alpha \bar{R}_{tb}^a \omega^b$. The vector field $\omega^\alpha = \omega^a e_{(a)}^\alpha$ was already introduced in Eq. (2.31), and its geometrical meaning was described at the end of Sec. II. The vector field $\bar{a}^a e_{(a)}^\alpha$ can easily be related to a^α , the acceleration field of Eq. (2.28); by combining Eqs. (2.15), (2.16), and (2.28) we obtain

$$\bar{a}^a e_{(a)}^\alpha = (1 + s\bar{a}_c \omega^c) a^\alpha - s e_{(a)}^\alpha \bar{R}_{tct}^a \omega^c + O(s^2). \quad (3.10)$$

Making this substitution into Eq. (3.9) and involving Eq. (3.3), we obtain

$$\begin{aligned} f^\alpha = e\bar{e} & \left[\frac{1}{s^2} \omega^\alpha - \frac{1}{2s} a^\alpha - \frac{1}{2s} (\bar{a}_b \omega^b) \omega^\alpha + \frac{1}{4} (\bar{a}_b \omega^b) a^\alpha \right. \\ & + \frac{3}{8} (\bar{a}_b \omega^b)^2 \omega^\alpha + \frac{3}{8} \dot{a}_t \omega^\alpha - \frac{1}{6} e_{(a)}^\alpha \bar{R}_{tbt}^a \omega^b \\ & - \frac{1}{6} \bar{R}_{tct} \omega^b \omega^c \omega^\alpha + \frac{1}{12} (5\bar{R}_{tt} + \bar{R}_{bc} \omega^b \omega^c + \bar{R}) \omega^\alpha \\ & \left. - \frac{1}{6} e_{(a)}^\alpha \bar{R}_{tb}^a \omega^b \right] + e e_{(a)}^\alpha F_{R(0)}^{(a)} + O(s). \end{aligned} \quad (3.11)$$

This expression for the force involves the vectors ω^α , a^α , and $e_{(a)}^\alpha$ which are all appropriately defined at $x = (t, s\omega^a)$, where the charge e resides. But it also involves quantities such as $\bar{a}_a$, $\dot{a}_t$, $\bar{R}_{tct}^a$, etc., which are defined instead at $\bar{x}$. The result is cumbersome, and our expression for the force needs to be rationalized. We shall therefore endeavor to express f^α entirely in terms of quantities that are defined at x .

To see how this can be achieved we present the details of a representative piece of the calculation: We shall show that the quantity $\bar{a}_a \omega^a$ that appears in Eq. (3.11) can be expressed as

$$\bar{a}_a \omega^a = a_\alpha \omega^\alpha + s(a_\alpha \omega^\alpha)^2 - s R_{\alpha\mu\beta\nu} \omega^\alpha u^\mu \omega^\beta u^\nu + O(s^2), \quad (3.12)$$

in terms of tensorial quantities defined at x only. Here a^α is the acceleration field of Eq. (2.28), u^α is the velocity field of Eqs. (2.24) and (2.25), and $R_{\alpha\mu\beta\nu}$ is the Riemann tensor evaluated at x .

We begin by stating that by virtue of Eq. (2.31) and the orthonormality of the tetrad vectors, $\bar{a}^a \omega_a$ is equal to $\bar{a}^a e_{(a)}^\alpha \omega_\alpha$. Combining this statement with Eq. (3.10) gives

$$\bar{a}_a \omega^a = (1 + s\bar{a}_c \omega^c) a_\alpha \omega^\alpha - s \omega_\alpha e_{(a)}^\alpha \bar{R}_{tct}^a \omega^c + O(s^2),$$

or

$$\bar{a}_a \omega^a = a_\alpha \omega^\alpha + s(a_\alpha \omega^\alpha)^2 - s \omega_\alpha e_{(a)}^\alpha \bar{R}_{tct}^a \omega^c + O(s^2).$$

By involving the tetrad vectors of Eqs. (2.8)–(2.16), the components of the Riemann tensor in FNC can be expressed in covariant form as

$$\bar{R}_{tct}^a = R_{\bar{\lambda}\bar{\nu}\bar{\rho}}^{\bar{\mu}} e_{\bar{\mu}}^{(a)} e_{(0)}^{\bar{\lambda}} e_{(c)}^{\bar{\nu}} e_{(0)}^{\bar{\rho}}.$$

The right-hand side is a scalar quantity, and this scalar can be evaluated at x instead of $\bar{x}$ at the cost of introducing a correction term of order s . We then have

$$\bar{R}_{tct}^a = R_{\lambda\nu\rho}^\mu e_{(a)}^\lambda e_{(0)}^\nu e_{(0)}^\rho + O(s),$$

and it follows that

$$e_{(a)}^\alpha \bar{R}_{tct}^a \omega^c = R_{\lambda\nu\rho}^\mu (e_{(a)}^\alpha e_{(0)}^\lambda e_{(c)}^\nu e_{(0)}^\rho) \omega^c + O(s).$$

The factor within the first set of brackets is $\delta_{\mu}^{\alpha} - e_{(0)}^{\alpha} e_{\mu}^{(0)}$, and the factor within the second set is ω^ν ; the right-hand side becomes

$$R_{\lambda\nu\rho}^\alpha e_{(0)}^\lambda \omega^\nu e_{(0)}^\rho + O(s).$$

We now compare Eqs. (2.13), (2.14), and (2.24), (2.25) and observe that $e_{(0)}^\alpha = u^\alpha + O(s^2)$. This allows us to write, finally,

$$e_{(a)}^\alpha \bar{R}_{tct}^a \omega^c = R_{\lambda\nu\rho}^\alpha u^\lambda \omega^\nu u^\rho + O(s),$$

and we have obtained Eq. (3.12).

Working on each term of Eq. (3.11) in this fashion, we eventually obtain our final expression

$$\begin{aligned} f^\alpha = & e\bar{e} \left[\frac{1}{s^2} \omega^\alpha - \frac{1}{2s} a^\alpha - \frac{1}{2s} (a_\beta \omega^\beta) \omega^\alpha + \frac{1}{4} (a_\beta \omega^\beta) a^\alpha \right. \\ & - \frac{1}{8} (a_\beta \omega^\beta)^2 \omega^\alpha + \frac{3}{8} (\dot{a}_\beta u^\beta) \omega^\alpha \\ & + \frac{1}{3} \omega^\alpha R_{\beta\gamma\delta\epsilon} \omega^\beta u^\gamma \omega^\delta u^\epsilon - \frac{1}{6} R_{\beta\gamma\delta}^\alpha u^\beta \omega^\gamma u^\delta \\ & + \frac{1}{12} (5R_{\beta\gamma} u^\beta u^\gamma + R_{\beta\gamma} \omega^\beta \omega^\gamma + R) \omega^\alpha \\ & \left. - \frac{1}{6} (\delta_\beta^\alpha + u^\alpha u_\beta) R_{\gamma}^\beta \omega^\gamma \right] + eF_{R\beta}^\alpha u^\beta \\ & + O(s) \end{aligned} \quad (3.13)$$

for the force acting on the charge e at position x , due to the charge $\bar{e}$ moving on $\bar{\gamma}$. This expression involves s , the spatial distance from $\bar{x}$ to x ; the unit vector ω^α defined at the end of Sec. II; the velocity vector u^α of the charge e , its acceleration vector a^α , and its covariant derivative $\dot{a}^\alpha$; the Riemann and Ricci tensors, and the Ricci scalar, all evaluated at x . And it involves the Detweiler-Whiting regular field, also evaluated at x [Eqs. (3.5)–(3.8); LRR Eqs. (473) and (474)],

$$\begin{aligned} eF_{R\beta}^\alpha u^\beta = & e\bar{e} \left[(\delta_\beta^\alpha + u^\alpha u_\beta) \left(\frac{2}{3} \dot{a}^\beta + \frac{1}{3} R_{\gamma}^\beta u^\gamma \right) \right. \\ & \left. + 2u_\beta \int_{-\infty}^{\bar{\tau}^-} \nabla^{[\alpha} G_{\gamma'}^{(\beta]}(x, z') u^{\gamma'} d\bar{\tau}' + O(s) \right]. \end{aligned} \quad (3.14)$$

It is important to note that our final expression for f^α is a legitimate tensorial relation that is valid in all coordinate systems; all traces of the Fermi normal coordinates, which were heavily involved in the calculations leading to Eq. (3.13), have been eliminated.

IV. DUMBBELL

We construct an extended charged body, a dumbbell, by placing a charge e_- on a accelerated world line γ_- , and by placing a second charge e_+ on a neighboring world line γ_+ . The magnitude of each charge is arbitrary, but we assume that their sum $e = e_- + e_+$, the total charge of the extended body, does not vanish. We imagine that each charge follows a predetermined flight plan, and we wish to determine the forces acting on each charge. In a suitable sense to be spelled out in the following section, the sum total of these forces will be the net force acting on the extended body.

We imagine that each charge is equipped with a control system that consists of a sensor capable of detecting any deviation from the charge's prescribed motion, and a propulsion system capable of applying the necessary corrections to return the charge to its prescribed world line.³ The acceleration of each world line is produced by an external force f_{ext}^α which varies smoothly from γ_- to γ_+ ; the acceleration of each world line is related to the external force by $ma^\alpha = f_{\text{ext}}^\alpha$, where m is the mass of each charge. The role of the propulsion system is to apply a force f_{prop}^α that compensates for the electromagnetic forces produced within the dumbbell, which tend to push each charge away from its prescribed motion. The sum total of the propulsion forces is equal and opposite to the sum total of the forces produced within the dumbbell.

The motion of each charge will be referred to a reference world line $\bar{\gamma}$ which defines the spatial origin of a Fermi coordinate system (t, x^a) . The reference world line lies between the charges, and this new Fermi system is *distinct* from the one used in the preceding section. The previous system of FNC was centered on one of the two charges, the one that was labeled $\bar{e}$; here the FNC are centered instead on the central world line $\bar{\gamma}$. Nevertheless, our construction will allow us to continue to use the language and results of Sec. II.

Because our charges are made to follow a predetermined flight plan, we are free to choose their world lines $\gamma_\pm$. We pick two members of the congruence of comoving world lines, and we pick them to be on opposite sides of the reference world line; this translates into the selection $\gamma_\pm = \gamma(x_\pm^a)$, where $x_\pm^a = \pm \varepsilon n^a$. Here, ε is the constant spatial separation between each world line and $\bar{\gamma}$, and n^a is a unit vector that specifies the orientation of the dumbbell. The charges are therefore maintained at a constant

³ Here we deviate from the Ori-Rosenthal treatment, in which the charges are maintained in place by a rigid rod instead of a propulsion system. We use the artifice of a propulsion system to avoid the artifice of a perfectly rigid rod (which would violate relativistic causality) or the complications (such as energy dissipation) associated with a physically realistic rod. We do not think that this difference in modeling is very important; in flat spacetime the two models produce identical answers for the self-force.

spatial separation $s = 2\varepsilon$.

The components of each charge's velocity vector in FNC can be obtained from Eqs. (2.24), (2.25) by substituting $x^a = x_\pm^a = \pm \varepsilon n^a$. This gives

$$u_\pm^t = 1 \mp \varepsilon \bar{a}_a n^a + \varepsilon^2 (\bar{a}_a n^a)^2 - \frac{1}{2} \varepsilon^2 \bar{R}_{ctdt} n^c n^d + O(\varepsilon^3), \quad (4.1)$$

$$u_\pm^a = 0, \quad (4.2)$$

where, as before, $\bar{a}_a$ are the components of $\bar{\gamma}$'s acceleration vector, and $\bar{R}_{ctdt}$ are components of the Riemann tensor evaluated on $\bar{\gamma}$. The acceleration vectors are given by Eq. (2.28),

$$a_\pm^t = O(\varepsilon^2), \quad (4.3)$$

$$a_\pm^a = (1 \mp \varepsilon \bar{a}_b n^b) \bar{a}^a \pm \varepsilon \bar{R}_{tct}^a n^c + O(\varepsilon^2). \quad (4.4)$$

The force f_+^α acting on the charge e_+ , due to the charge e_- , can be obtained from Eq. (3.13) by making the substitutions $e = e_+$, $\bar{e} = e_-$, $s = 2\varepsilon$, $\omega^t = 0$, $\omega^a = +n^a$, $u^\alpha = u_+^\alpha$, $a^\alpha = a_+^\alpha$, and by involving Eqs. (4.1)–(4.4). Similarly, the force f_-^α acting on e_- , due to e_+ , can be obtained by making the substitutions $e = e_-$, $\bar{e} = e_+$, $s = 2\varepsilon$, $\omega^t = 0$, $\omega^a = -n^a$, $u^\alpha = u_-^\alpha$, and $a^\alpha = a_-^\alpha$. Straightforward manipulations reveal that the spatial components of the forces $f_\pm^\alpha$ in FNC are

$$\begin{aligned} f_\pm^a = e_+ e_- & \left[\pm \frac{1}{4\varepsilon^2} n^a - \frac{1}{4\varepsilon} \bar{a}^a - \frac{1}{4\varepsilon} (\bar{a}_b n^b) n^a \right. \\ & \pm \frac{1}{2} (\bar{a}_b n^b) \bar{a}^a \pm \frac{1}{8} (\bar{a}_b n^b)^2 n^a \pm \frac{3}{8} \dot{\bar{a}}_t n^a \\ & \pm \frac{1}{12} n^a \bar{R}_{btct} n^b n^c \mp \frac{5}{12} \bar{R}_{tbt}^a n^b \\ & \pm \frac{1}{12} (5\bar{R}_{tt} + \bar{R}_{bc} n^b n^c + \bar{R}) n^a \\ & \left. \mp \frac{1}{6} \bar{R}_{ab} n^b + \bar{f}_R^a + O(s) \right]. \end{aligned} \quad (4.5)$$

The time component of the forces is $O(s)$. In the manipulations leading to Eq. (4.5) we have displaced the evaluation point of the Riemann and Ricci tensors from $\gamma_\pm$ to the reference world line $\bar{\gamma}$. This can be done carefully by going through the procedure outlined in the paragraph following Eq. (3.12); or this can be done more simply by noting that the displacement introduces a correction term of order s that can be absorbed into the error term, $O(s)$.

We have introduced the rescaled Detweiler-Whiting regular force $\bar{f}_R^\alpha$, which is also evaluated on $\bar{\gamma}$ in Eq. (4.5). This is defined by removing a factor of the charge from the expression for $F_{R\beta}^\alpha u^\beta$ given in Eq. (3.14). If, for example, $e_+ F_{R\beta}^\alpha u_+^\beta$ is the contribution to the force acting on e_+ that comes from e_- 's regular field, we write it as $e_+ e_- \bar{f}_R^\alpha$ and displace the result from γ_+ to $\bar{\gamma}$. The rescaled regular force evaluated on $\bar{\gamma}$ is expressed in covariant form as

$$\bar{f}_R^\alpha = (\delta_{\bar{\beta}}^\alpha + u^{\bar{\alpha}} u_{\bar{\beta}}) \left(\frac{2}{3} \dot{\bar{a}}^{\bar{\beta}} + \frac{1}{3} R_{\bar{\gamma}}^{\bar{\beta}} u^{\bar{\gamma}} \right)$$

$$+ 2u_{\bar{\beta}} \int_{-\infty}^{\bar{\tau}^-} \nabla^{[\bar{\alpha}} G^{\bar{\beta}]}_{\bar{\gamma}'}(\bar{x}, z') u^{\bar{\gamma}'} d\bar{\tau}'. \quad (4.6)$$

It is independent of the charges $e_\pm$, and its spatial components in FNC are $\bar{f}_R^a$, which is what appears in Eq. (4.5).

V. TOTAL FORCE ACTING ON THE EXTENDED BODY

The net force acting on the dumbbell of Sec. IV consists of eight separate contributions. The first is f_+^a , the force acting on e_+ due to the field created by e_- ; the second is f_-^a , the force acting on e_- due to the field created by e_+ ; the third is $f_{\text{self}+}^a$, the self-force acting on e_+ due to this charge's own field; the fourth is $f_{\text{self}-}^a$, the self-force acting on e_- ; the fifth is $f_{\text{prop}+}^a$, the force produced by e_+ 's propulsion system to compensate for the action of f_+^a and $f_{\text{self}+}^a$; the sixth is $f_{\text{prop}-}^a$, the force produced by e_- 's propulsion system; and the remaining two are the external forces f_{ext}^a that are responsible for the acceleration of each charge.

We recall from Sec. I the fundamental axiom behind our derivation: *The self-force acting on a point charge is a well-defined vector field on the particle's world line, and it is proportional to the square of the particle's electric charge.* We will invoke these properties in the following manipulations, but we state now that according to the axiom, the individual self-forces $f_{\text{self}\pm}^a$ *must be included* in the accounting of the net force; it would be inconsistent to leave them out, as we expect the extended body to retain a nonvanishing self-force in the limit $\varepsilon \rightarrow 0$.

In a sense to be specified below, the net force acting on the dumbbell must be the sum total of the eight individual contributions listed above. We denote by $f_{\text{net,int}}^\alpha$ the net internal electromagnetic force acting on the dumbbell; this is the sum total of f_+^a , f_-^a , $f_{\text{self}+}^a$, and $f_{\text{self}-}^a$ evaluated on the central world line. We denote by $f_{\text{net,prop}}^\alpha$ the net force produced by the propulsion system of each charge; this is the sum total of $f_{\text{prop}+}^a$ and $f_{\text{prop}-}^a$ evaluated on the central world line. Finally, we denote by f_{ext}^α the net external force acting on the dumbbell; this is the sum total of the external forces acting on each charge.

The propulsion forces are designed to compensate for the action of the internal electromagnetic forces. We therefore have $f_{\text{net,prop}}^\alpha + f_{\text{net,int}}^\alpha = 0$, and the equation of motion for the dumbbell is $ma^\alpha = f_{\text{ext}}^\alpha$, where m is the total mass of the dumbbell; the external forces are keeping the dumbbell on its prescribed world line $\bar{\gamma}$. If we imagine ourselves momentarily switching off the propulsion system, the internal electromagnetic forces would no longer be compensated for, and we would expect that the dumbbell would start moving according to the new equation $ma^\alpha = f_{\text{ext}}^\alpha + f_{\text{self}}^\alpha$, with the second term denoting the self-force acting on the dumbbell.⁴ It is therefore

⁴ This equation is meant to apply only at the instant at which

clear that the propulsion forces are in fact compensating for the self-force, so that $f_{\text{net,prop}}^\alpha = -f_{\text{self}}^\alpha$. This allows us to conclude that the self-force acting on the dumbbell is equal to $f_{\text{net,int}}^\alpha$, and this is what we shall calculate in the remainder of this section.

The individual forces that make up $f_{\text{net,int}}^\alpha$ cannot simply be added up, because they are vectors that are defined at different points in the manifold of a curved spacetime. Before taking the sum they must all be transported to a single common point, which we choose to be $\bar{x}$ on the reference world line. We need a *transport rule* to carry out this operation.

The correct transport rule in flat spacetime was identified by Ori and Rosenthal [9, 10]. Building on considerations of momentum conservation, they showed that the transport rule involves parallel transport of the force from $\gamma_\pm$ to $\bar{\gamma}$ combined with multiplication by a kinematic factor $d\tau_\pm/d\bar{\tau}$ that converts increments of proper time on $\gamma_\pm$ to an increment of proper time on the central world line. The parallel-transport component of the rule is easy enough to understand, and the kinematic conversion is easily justified by the fact that an increment of momentum on each world line is related to the force by $dp_\pm^\alpha = f_\pm^\alpha d\tau_\pm$; because a similar statement applies to $\bar{\gamma}$, a conversion from $d\tau_\pm$ to $d\bar{\tau}$ is clearly required.

We write the net internal electromagnetic force acting on the dumbbell as

$$f_{\text{net,int}}^\alpha = k_\alpha^{\bar{\alpha}}(f_+^\alpha + f_{\text{self}+}^\alpha) + k_\alpha^{\bar{\alpha}}(f_-^\alpha + f_{\text{self}-}^\alpha), \quad (5.1)$$

where $k_\alpha^{\bar{\alpha}}(\bar{x}, x)$ is the transport operator which takes a force at $x_\pm$ on $\gamma_\pm$ and brings it to the point $\bar{x}$ on $\bar{\gamma}$. According to Ori and Rosenthal, this is $k_\alpha^{\bar{\alpha}} = (d\tau_\pm/d\bar{\tau})g_\alpha^{\bar{\alpha}}$ in flat spacetime, with the second factor denoting parallel transport from $x_\pm$ to $\bar{x}$. According to Eqs. (2.22) and (2.23), from which we remove all terms that involve the Riemann tensor, the relevant components of the transport operator in FNC are

$$k_b^a = (1 \pm \varepsilon \bar{a}_c n^c) \delta_b^a + O(\varepsilon^3) \quad (\text{flat spacetime}). \quad (5.2)$$

In curved spacetime we would expect this expression to be modified by a term of order ε^2 which, presumably, would be constructed from the Riemann tensor. We shall, however, remain uncommitted as to the exact nature of the transport rule in curved spacetime. We shall not assume that the flat-spacetime prescription $k_\alpha^{\bar{\alpha}} = (d\tau_\pm/d\bar{\tau})g_\alpha^{\bar{\alpha}}$ is preserved in curved spacetime. And we shall not adopt any specific alternative prescription such as, for example, the Dixon-Bailey-Israel transport rule [14, 15, 16, 17, 18] that is widely used in the description of extended bodies in general relativity. Instead, we shall adopt a very generic transport rule of the form

$$k_b^a = (1 \pm \varepsilon \bar{a}_c n^c) \delta_b^a + \varepsilon^2 \bar{A}_{bcd} n^c n^d + O(\varepsilon^3), \quad (5.3)$$

with $\bar{A}_{bcd}^a$ representing the components of an unknown tensor which is evaluated at $\bar{x}$. We may imagine that this tensor is formed from the Riemann tensor, but we do not need to make such a statement; our only requirement here is that the quantities $\bar{A}_{bcd}^a$ must depend on $\bar{x}$ only and be independent of $x_\pm$. The only allowed dependence of k_b^a on $x_\pm = \pm \varepsilon n^a$ is the one which is shown explicitly in Eq. (5.3).

Observe that Eq. (5.3) can be viewed as representing the Taylor series of a general bitensor k_b^a in powers of x^a . The first two terms in the series are determined by the requirement that k_b^a must reduce to the expression of Eq. (5.2) in flat spacetime. The term of order ε^2 , however, is arbitrary. Its only relevant property, for the purposes of our calculation, is that *it comes with the same sign* whether the transport originates from x_- or x_+ ; unlike the term of order ε , it does not alternate in sign.

Combining Eqs. (4.5) and (5.3) gives

$$\begin{aligned} k_b^a f_\pm^b = & e_+ e_- \left[\pm \frac{1}{4\varepsilon^2} n^a - \frac{1}{4\varepsilon} \bar{a}^a \pm \frac{1}{4} (\bar{a}_b n^b) \bar{a}^a \right. \\ & \mp \frac{1}{8} (\bar{a}_b n^b)^2 n^a \pm \frac{3}{8} \bar{a}_t n^a \pm \frac{1}{12} n^a \bar{R}_{btct} n^b n^c \\ & \mp \frac{5}{12} \bar{R}_{tbt} n^b \pm \frac{1}{12} (5\bar{R}_{tt} + \bar{R}_{bc} n^b n^c + \bar{R}) n^a \\ & \mp \frac{1}{6} \bar{R}_b^a n^b \pm \frac{1}{4} \bar{A}_{bcd}^a n^b n^c n^d \\ & \left. + \bar{f}_R^a + O(\varepsilon) \right], \end{aligned} \quad (5.4)$$

and from this it follows that

$$k_b^a f_+^b + k_b^a f_-^b = e_+ e_- \left[-\frac{1}{2\varepsilon} \bar{a}^a + 2\bar{f}_R^a + O(\varepsilon) \right]. \quad (5.5)$$

This result will be substituted into Eq. (5.1) in a moment.

The net internal electromagnetic force acting on the dumbbell involves also the combined action of the individual self-forces $f_{\text{self}\pm}^a$. Here we invoke our axiom and write

$$f_{\text{self}\pm}^a = e_\pm^2 f_{\text{self}}^a[\gamma_\pm], \quad (5.6)$$

which makes the statement that each self-force is proportional to the square of the particle's charge; the rescaled self-force $f_{\text{self}}^a[\gamma_\pm]$ is independent of $e_\pm$. As indicated, each rescaled self-force is a functional of a world line which, we presume, can be defined in terms of the velocity vector and its derivatives, and in terms of spacetime tensors (and bitensors) that are evaluated on the world line. We assume that this is *the same functional* for each world line, and that the result is a smooth vector field on the world line. From Eqs. (5.3) and (5.6) we get

$$k_b^a f_{\text{self}+}^b + k_b^a f_{\text{self}-}^b = (e_+^2 + e_-^2) \bar{f}_{\text{self}}^a[\bar{\gamma}] + O(\varepsilon), \quad (5.7)$$

in which the rescaled self-force is now evaluated on the central world line $\bar{\gamma}$.

the propulsion system is switched off. We need not be concerned with the fact that the dumbbell has momentarily lost its cohesion and is about to fly apart.

Equation (5.1) now reads

$$\begin{aligned} \bar{f}_{\text{net,int}}^a &= -\frac{e_+e_-}{s}\bar{a}^a + 2e_+e_-\bar{f}_R^a + (e_+^2 + e_-^2)\bar{f}_{\text{self}}^a[\bar{\gamma}] \\ &\quad + O(s), \end{aligned} \quad (5.8)$$

where $s = 2\varepsilon$ is the spatial separation between the two charges. Equation (5.8) gives the components of the net force in FNC.

The first term on the right-hand side of Eq. (5.8) is divergent in the limit $s \rightarrow 0$. It has the form of $-m_{\text{el}}a^a$, where

$$m_{\text{el}} = \frac{e_+e_-}{s} \quad (5.9)$$

is the electromagnetic contribution — its electrostatic energy — to the mass of the dumbbell. This term produces an additional inertial term in the equations of motion, and it can be combined with the term $m_{\text{material}}a^a$ to form the appropriate left-hand side, $m_{\text{inertial}}a^a = (m_{\text{material}} + m_{\text{el}})a^a$. Here m_{material} is the purely material contribution to the dumbbell's mass, to which the electromagnetic contribution m_{el} is added to form m_{inertial} , the dumbbell's total inertial mass. The diverging term has therefore been eliminated by mass renormalization, and at this stage there is no obstacle in taking the limit $s \rightarrow 0$ of Eq. (5.8); the dumbbell finally becomes a point particle.

As was first pointed out by Ori and Rosenthal [9, 10], the mass-renormalization procedure is successful because the diverging term in $\bar{f}_{\text{net}}^a$ is precisely proportional to the acceleration $\bar{a}^a$; this is a true property of the net force because the directional terms $(\bar{a}_b n^b)n^a/(4\varepsilon)$ that appeared in the mutual forces $f_{\pm}^a$ were eliminated during the transport from $\gamma_{\pm}$ to $\bar{\gamma}$. As they also pointed out, the procedure is successful also because it is not subjected to the famous 4/3 problem [19]: the shift in the mass is m_{el} and not $\frac{4}{3}m_{\text{el}}$.

The second and third terms in Eq. (5.8) combine to form the dumbbell's own self-force,

$$\bar{f}_{\text{self}}^a = 2e_+e_-\bar{f}_R^a + (e_+^2 + e_-^2)\bar{f}_{\text{self}}^a[\bar{\gamma}]. \quad (5.10)$$

We once more invoke our axiom and write the self-force as

$$\bar{f}_{\text{self}}^a = e^2\bar{f}_{\text{self}}^a[\bar{\gamma}], \quad (5.11)$$

in terms of the total charge $e = e_+ + e_-$ and in terms of the same rescaled self-force functional that was introduced in Eq. (5.6). Equating Eqs. (5.10) and (5.11) produces an equation to be solved for $\bar{f}_{\text{self}}^a[\bar{\gamma}]$, and we obtain our central result,

$$\bar{f}_{\text{self}}^a[\bar{\gamma}] = \bar{f}_R^a. \quad (5.12)$$

The rescaled self-force is the Detweiler-Whiting rescaled regular force of Eq. (4.6).

Our final conclusion is that the self-force acting on a point particle of charge e is given by

$$\begin{aligned} f_{\text{self}}^{\bar{\alpha}} &= e^2(\delta^{\bar{\alpha}}_{\bar{\beta}} + u^{\bar{\alpha}}u_{\bar{\beta}})\left(\frac{2}{3}\dot{a}^{\bar{\beta}} + \frac{1}{3}R^{\bar{\beta}}_{\bar{\gamma}}u^{\bar{\gamma}}\right) \\ &\quad + 2e^2u_{\bar{\beta}}\int_{-\infty}^{\bar{\tau}^-}\nabla^{[\bar{\alpha}}G^{\bar{\beta}]}_{\bar{\gamma}'}(\bar{x}, z')u^{\bar{\gamma}'}d\bar{\tau}'. \end{aligned} \quad (5.13)$$

This is the standard expression for the self-force. The particle's equations of motion are then

$$ma^{\bar{\alpha}} = f_{\text{self}}^{\bar{\alpha}} + f_{\text{ext}}^{\bar{\alpha}}, \quad (5.14)$$

with the inertial mass m identified with the combined material and electromagnetic mass of the charged body.

VI. DISCUSSION: CRITICAL REVIEW OF EXISTING DERIVATIONS OF THE SELF-FORCE

We review in this section the various approaches that have been used in the literature to derive the standard expression for the electromagnetic self-force in curved spacetime, as given by Eq. (1.2). While all these approaches have led to the same answer, they possess different strengths and weaknesses, and none of them are completely immune to criticism. This critical review motivated our own approach to the problem, which is also summarized at the end of this section. The presentation in this section borrows heavily from [LRR Sec. 5.5].

A. Conservation of energy-momentum

Conservation of energy-momentum was used by Dirac [2] to derive the equations of motion of a point electric charge in flat spacetime, and the same method was adopted by DeWitt and Brehme [1] in their generalization of Dirac's work to curved spacetime. This approach, of course, produced the first derivation of the standard expression. (The DeWitt-Brehme derivation contained a calculational mistake that was later corrected by Hobbs [3].)

The method is based on the conservation equation $T^{\alpha\beta}_{;\beta} = 0$, where the stress-energy tensor $T^{\alpha\beta}$ includes a contribution from the charged particle and a contribution from the electromagnetic field. The particle's contribution is a Dirac distribution on the world line, and the field's contribution diverges as $1/s^4$ near the world line, where s is the spatial distance from the field point to the world line. (We are using Fermi normal coordinates in this discussion.) While in flat spacetime the differential statement of energy-momentum conservation can immediately be turned into an integral statement, the same is not true in a curved spacetime (unless the spacetime possesses at least one Killing vector). To proceed it is necessary to rewrite the conservation equation as

$$0 = g^{\bar{\mu}}_{\bar{\alpha}}T^{\alpha\beta}_{;\beta} = (g^{\bar{\mu}}_{\bar{\alpha}}T^{\alpha\beta})_{;\beta} - g^{\bar{\mu}}_{\alpha;\beta}T^{\alpha\beta}, \quad (6.1)$$

where $g_{\alpha}^{\bar{\mu}}(\bar{x}, x)$ is the parallel propagator from x to an arbitrary point $\bar{x}$ on the world line. Integrating this equation over the interior of a world-tube segment that consists of a “wall” of constant s and two “caps” of constant t , we obtain

$$0 = \int_{\text{wall}} g_{\alpha}^{\bar{\mu}} T^{\alpha\beta} d\Sigma_{\beta} + \int_{\text{caps}} g_{\alpha}^{\bar{\mu}} T^{\alpha\beta} d\Sigma_{\beta} + \int_{\text{interior}} g_{\alpha;\beta}^{\bar{\mu}} T^{\alpha\beta} dV, \quad (6.2)$$

where $d\Sigma_{\beta}$ is a three-dimensional surface element and dV an invariant, four-dimensional volume element.

There is no obstacle in evaluating the wall integral, for which $T^{\alpha\beta}$ reduces to the field’s stress-energy tensor; for a wall of radius s the integral scales as $1/s^2$. The integrations over the caps, however, are problematic: While the particle’s contribution to the stress-energy tensor is integrable, the integration over the field’s contribution goes as $\int_0^s (s')^{-2} ds'$ and diverges. This issue arises also in flat spacetime [2], and in this case the regularization of the cap integrations, and the removal of all singular terms by mass renormalization, has been done with great rigor using robust distributional methods. The state of the art for flat spacetime was reviewed in a paper by Teitelboim, Villarroel, and van Weert [20]. As far as we are aware, however, the cap integrations for curved spacetime were never handled with such care, and the DeWitt-Brehme derivation of the self-force does not enjoy the same degree of rigor as what has been achieved for flat spacetime.

More troublesome, however, is the treatment of the interior integral, which does not occur in flat spacetime, and which was simply discarded by DeWitt and Brehme. Because $g_{\alpha;\beta}^{\bar{\mu}}$ scales as s , the field’s stress-energy tensor as $1/s^4$, and dV as $s^2 ds dt$, this integral goes as $\int_0^s (s')^{-1} ds' dt$ and it also diverges, albeit less strongly than the caps integration. While simply discarding this integral does produce the standard expression for the electromagnetic self-force, it would be desirable to go through a careful regularization of the interior integration, and find a convincing reason to discard it altogether. To the best of our knowledge, this has not been done, and until this issue is fully resolved the DeWitt-Brehme derivation shall be open to criticism.

B. Averaging method

An alternative derivation of the electromagnetic self-force is presented in [LRR Sec. 5.2.6]. It is based on the idea that it is the average of the retarded field around the charged particle which is responsible for the net force acting on it. This derivation has the advantage of being much easier to implement than a derivation based on energy-momentum conservation; it requires a computation of the retarded field of a point charge, but it does not require the computation of the field’s stress-energy tensor.

To describe the averaging method it is convenient to adopt the Detweiler-Whiting decomposition of the retarded field $F_{\alpha\beta}$ into singular and regular pieces [5],

$$F_{\alpha\beta} = F_{\alpha\beta}^S + F_{\alpha\beta}^R. \quad (6.3)$$

As was shown by Detweiler and Whiting, this decomposition is unambiguous, the retarded and singular fields share the same singularity structure near the world line, and the regular field $F_{\alpha\beta}^R$ is smooth on the world line.

To formulate equations of motion for the point charge we temporarily model it as a spherical hollow shell, and we obtain the net force acting on this object by averaging $F_{\alpha\beta}$ over the shell’s surface. (The averaging is performed in the shell’s rest frame, and the shell is spherical in the sense that its proper distance s from the world line is the same in all directions.) The averaged field is next evaluated on the world line, in the limit of a zero-radius shell. Because the regular field is smooth on the world line, this yields

$$e\langle F_{\bar{\mu}\bar{\nu}} \rangle u^{\bar{\nu}} = e\langle F_{\bar{\mu}\bar{\nu}}^S \rangle u^{\bar{\nu}} + eF_{\bar{\mu}\bar{\nu}}^R u^{\bar{\nu}}, \quad (6.4)$$

where $u^{\bar{\mu}}$ is the particle’s velocity vector. As was calculated in [LRR Eq. (475)],

$$e\langle F_{\bar{\mu}\bar{\nu}}^S \rangle u^{\bar{\nu}} = -(m'_{\text{el}})a_{\bar{\mu}}, \quad m'_{\text{el}} = \frac{2}{3} \frac{e^2}{s}, \quad (6.5)$$

and $eF_{\bar{\mu}\bar{\nu}}^R u^{\bar{\nu}}$ was given in Eq. (5.13). The equations of motion are then postulated to be $m_{\text{material}} a_{\bar{\mu}} = e\langle F_{\bar{\mu}\bar{\nu}} \rangle u^{\bar{\nu}} + f_{\bar{\mu}}^{\text{ext}}$, where m_{material} is the particle’s material mass. With the preceding results we arrive at $m_{\text{inertial}} a_{\bar{\mu}} = eF_{\bar{\mu}\bar{\nu}}^R u^{\bar{\nu}} + f_{\bar{\mu}}^{\text{ext}}$, where $m_{\text{inertial}} \equiv m_{\text{material}} + m'_{\text{el}}$ is the particle’s observed (renormalized) inertial mass. We have recovered the standard expression for the self-force, $f_{\bar{\mu}}^{\text{self}} = eF_{\bar{\mu}\bar{\nu}}^R u^{\bar{\nu}}$.

The averaging method is basically sound, but it contains a number of weak points. A first source of criticism concerns the specifics of the averaging procedure, in particular, the choice of a spherical surface over any other conceivable shape. Another source is an inconsistency of the method which gives rise to the famous “4/3 problem” [19]: the mass shift m'_{el} is related to the shell’s electrostatic energy $m_{\text{el}} = e^2/(2s)$ by $m'_{\text{el}} = \frac{4}{3}m_{\text{el}}$ instead of the expected $m'_{\text{el}} = m_{\text{el}}$. This problem originates from the fact that the field that is averaged over the surface of the shell is sourced by a point particle and not by the shell itself. It is likely that a more careful treatment of the near-source field will eliminate both sources of criticism: We can expect that the field produced by an extended spherical object will give rise to a mass shift that equals the object’s electrostatic energy, and the object’s spherical shape would then fully justify a spherical averaging. (Considering other shapes might also be possible, but one would prefer to keep the object’s structure simple and avoid introducing additional multipole moments.) Further work is required to clean up these details.

The averaging method is at the core of the approach followed by Quinn and Wald [4], who also average the

retarded field over a spherical surface surrounding the particle. Their approach, however, also incorporates a “comparison axiom” that eliminates the need to explicitly renormalize the mass. Their axiomatic approach is perfectly sound, but it completely bypasses the interesting question of determining the electromagnetic contribution to the particle’s inertia.

C. Detweiler-Whiting axiom

The Detweiler-Whiting decomposition of the retarded field [5] becomes most powerful when it is combined with the Detweiler-Whiting axiom, which asserts that

the singular field $F_{\alpha\beta}^S$ exerts no force on the particle (it merely contributes to the particle’s inertia); the entire self-force arises from the action of the regular field $F_{\alpha\beta}^R$.

This axiom, which is motivated by the time-symmetric nature of the singular field and its causal relation with the world line, gives rise to the equations of motion $ma_{\bar{\mu}} = eF_{\bar{\mu}\bar{\nu}}^R u^{\bar{\nu}} + f_{\bar{\mu}}^{\text{ext}}$, in agreement with the averaging method (but with an implicit, instead of explicit, mass shift), and in agreement with the standard expression for the electromagnetic self-force. In this picture, the particle simply interacts with a free field whose origin can be traced to the particle’s past, and the procedure of mass renormalization is again sidestepped. This picture of a particle interacting with a free field is compelling, and it removes any tension between the nongeodesic motion of the charge and the principle of equivalence. But the Detweiler-Whiting axiom is quite strong, and one would like to justify it on the basis of an independent derivation of the self-force. This observation leads us toward an extended-body approach as described in Sec. I, and toward our particular partial implementation of this program.

D. Extended-body approach

The derivation of the electromagnetic self-force presented in this paper is a generalization to curved spacetime of the extended-body approach of Ori and Rosenthal [9, 10], which was limited to flat spacetime. It is an attempt to eliminate the difficulties associated with a point particle by dealing instead with an extended charge distribution. Our attempt is only partially successful: While we do introduce an extended body, it still consists of two point charges, and we make no attempt to deal with the internal dynamics, which we replace with a contrived and artificial control system that keeps the individual charges on their predetermined world lines. The result, to be sure, is a derivation of the standard expression for the self-force that lacks a strong physical backbone. But while we readily acknowledge this major flaw, we stress nevertheless that our derivation is

logically sound and consistent, and that it leads to the correct answer displayed in Eq. (1.2). Taking the view that all details regarding internal structure must become irrelevant in the limit $s \rightarrow 0$, perhaps it is not so objectionable that in a first attempt to describe the motion of an extended charged body in curved spacetime, one would adopt an internal structure that offers simplicity at the cost of realism.

Our extended body does indeed have the simplest conceivable shape: It is a dumbbell consisting of two point charges maintained at a constant spacelike separation s ; the dumbbell’s orientation relative to the direction of motion is arbitrary. The internal electromagnetic forces acting on this extended body can be broken down into four contributions: The first two are the mutual forces that each charge exerts on the other, and the remaining two are the self-forces of each individual charge. The self-force acting on the entire extended body is the sum total of these four contributions, combined appropriately in a way that respects the curved nature of the spacetime. To identify the proper way of adding the forces was a key aspect of our derivation; here we followed the Ori-Rosenthal prescription and generalized it to curved spacetime.

Another key aspect of this derivation was our reliance on the fact that *each point charge within the dumbbell is subjected to its own self-force*. That this is so may seem peculiar, but it is in fact required by physical consistency: It would not do to assume that the individual self-forces do not exist and to find that a nonzero self-force eventually emerges from the extended body in the limit $s \rightarrow 0$. A successful extended-body calculation based on a collection of point charges must therefore incorporate the idea that each charge is subjected to its own self-force.

The extended-body approach is a consistent method of derivation that yields the standard expression for the electromagnetic self-force. It does, however, require an input axiom, which we reproduce from Sec. I:

The self-force acting on a point charge is a well-defined vector field on the particle’s world line, and it is proportional to the square of the particle’s electric charge.

The fact that the self-force must exist and be a well-defined quantity was motivated in the preceding paragraph. The other important aspect of the axiom is the statement that the self-force must scale like the square of the charge. That this must be so follows again from a consistency requirement: In a spacetime with a timelike Killing vector, the total work done by the self-force must equal the total energy radiated by the particle [11]; because the radiated energy is proportional to the square of the charge, the same should be required of the self-force.

As was discussed in Sec. I, the need for such an axiom is unfortunate, and we must add this flaw to the flaws that have already been identified. Such an axiom would not be required in an extended-body calculation that would deal honestly with the body’s internal dynamics. This flaw, therefore, is very much a consequence

of our refusal to introduce a physical model for the extended charge distribution and the cohesive forces that hold it together. In addition, the assumed equality between radiated energy and work done by the self-force must be only approximately valid, because heat dissipation within an extended body would also contribute to the energy balance. This effect, however, should become unimportant in the limit $s \rightarrow 0$, that is, in the limit in which the internal dynamics decouple from the external dynamics.

We regard this axiom as a necessary evil, but we see it also as a fairly mild assumption. It is, in particular, milder than the Detweiler-Whiting axiom, and we may view the extended-body approach as eventually providing a means to relax the main assumption behind the Detweiler-Whiting approach. What we would have, then, is an independent justification of the statement that the singular field exerts no force on the particle, but that it makes an electrostatic-energy contribution to its inertial mass.

We have tried to present a fair and balanced assessment of the weaknesses and strengths of our dumbbell model. While our derivation of the electromagnetic self-force does produce the standard expression of Eq. (1.2), it leaves much to be desired in terms of physical realism. Our refusal to deal with the body's internal dynamics comes with a high price, and our model is indeed too ar-

tificial and contrived (though still consistent) to be fully satisfactory. It would be desirable to return to this subject with a more physical model for the extended charge distribution. We believe that the calculational tools introduced in this paper will be useful in this endeavor. In particular, it is clear to us that the problem is best formulated in terms of Fermi normal coordinates, which incorporate in a general and straightforward way the curved nature of the spacetime. Another valuable contribution is our demonstration, in Sec. V, that the addition of partial forces acting within the extended body is largely insensitive to the details of the transport rule that carries these forces to a common point. We hope that in the future, these techniques will facilitate the formulation of a more realistic internal model.

Acknowledgments

We thank Eran Rosenthal for useful discussions. We also thank an anonymous referee for very valuable comments on an earlier version of this article, in which the flaws of our approach were not so clearly identified. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] B. S. DeWitt and R. W. Brehme, *Ann. Phys. (N.Y.)* **9**, 220 (1960).
 - [2] P. A. M. Dirac, *Proc. Roy. Soc. London* **A167**, 148 (1938).
 - [3] J. M. Hobbs, *Ann. Phys. (N.Y.)* **47**, 141 (1968).
 - [4] T. C. Quinn and R. M. Wald, *Phys. Rev. D* **56**, 3381 (1997), arXiv:gr-qc/9610053.
 - [5] S. Detweiler and B. F. Whiting, *Phys. Rev. D* **67**, 024025 (2003), arXiv:gr-qc/0202086.
 - [6] E. Poisson, *Living Rev. Relativity* **7** (2004), 6. [Online article]: cited on February 7, 2008, <http://www.livingreviews.org/lrr-2004-6>.
 - [7] Y. Mino, M. Sasaki, and T. Tanaka, *Phys. Rev. D* **55**, 3457 (1997), arXiv:gr-qc/9712056.
 - [8] E. Poisson, *Phys. Rev. D* **69**, 084007 (2004), arXiv:gr-qc/0311026.
 - [9] A. Ori and E. Rosenthal, *Phys. Rev. D* **68**, 041701 (2003), arXiv:gr-qc/0205003.
 - [10] A. Ori and E. Rosenthal, *J. Math. Phys.* **45**, 2347 (2004), arXiv:gr-qc/0309102.
 - [11] T. C. Quinn and R. M. Wald, *Phys. Rev. D* **60**, 064009 (1999), arXiv:gr-qc/9903014.
 - [12] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
 - [13] E. Poisson, *A relativist's toolkit: The mathematics of black-hole mechanics* (Cambridge University Press, Cambridge, England, 2004).
 - [14] W. G. Dixon, *Proc. Roy. Soc. London* **A314**, 499 (1970).
 - [15] W. G. Dixon, *Proc. Roy. Soc. London* **A319**, 509 (1970).
 - [16] W. G. Dixon, *Phil. Trans. Roy. Soc. London* **A277**, 59 (1974).
 - [17] W. G. Dixon, in *Isolated systems in general relativity*, edited by J. Ehlers (North Holland, Amsterdam, Netherlands, 1979).
 - [18] I. Bailey and W. Israel, *Ann. Phys. (N.Y.)* **130**, 188 (1980).
 - [19] F. Rohrlich, *Classical charged particles* (Addison-Wesley, Redwood City, 1990).
 - [20] C. Teitelboim, D. Villarroel, and C. G. van Weert, *Riv. Nuovo Cimento* **3**, 9 (1980).